

ARKADY M. ALT

ABSTRACT. This paper aim is consideration of elementary (calculus-free) way which only the double-angle formulas and the inequalities $\sin x < x < \tan x$ give opportunity to obtain well known polynomial minorants and majorants for $\sin x$, $\cos x$ and $\tan x$.

1. Introduction

We offer an elementary (calculus-free) proof of the inequalities

$$x - \frac{x^3}{6} < \sin x < x - \frac{x^3}{6} + \frac{x^5}{120}, \quad (1)$$

$$1 - \frac{x^2}{2} < \cos x < 1 - \frac{x^2}{2} + \frac{x^4}{24}, \quad (2)$$

$$x + \frac{x^3}{3} < \tan x, \quad (3)$$

when $x \in \left(0, \frac{\pi}{2}\right)$. All we need are the double-angle formulas

$$\begin{aligned} \sin 2x &= 2 \sin x \cos x, \\ \cos 2x &= 1 - 2 \sin^2 x, \\ \tan 2x &= \frac{2 \tan x}{1 - \tan^2 x}, \end{aligned}$$

all presented in standard courses of trigonometry, and the double inequality $\sin x < x < \tan x$ for $x \in (0, \pi/2)$, which has a visual geometric proof.

The left side of (2) can be improved to become

$$1 - \frac{x^2}{2} < 1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} < \cos x \quad (4)$$

and we prove this as well. In each case the polynomials consist of the initial terms of Maclaurin series for the sine, cosine, and tangent functions.

When we say "calculus-free proof" we mean that proof should be free not only from derivative and series but also free from limits, using instead "passing to limit", the simple and transparent reasoning in form of following

Proposition 1. *Let P be set of positive real numbers such that for any positive real ε there is $p \in P$ that $p < \varepsilon$ and let inequality $f(x) < p$ holds for any $x \in \text{Dom}(f)$ and any $p \in P$. Then for any $x \in \text{Dom}(f)$ holds inequality $f(x) \leq 0$.*

Proof. Indeed, supposition of existence $x_0 \in \text{Dom}(f)$ such that $f(x_0) > 0$ immediately leads to contradiction because then $f(x_0) < p$ for any $p \in P$ and at the same time for $\varepsilon = f(x_0)$ there is $p \in P$ that $p < f(x_0)$. \square

• **Proof of $x - x^3/6 < \sin x$, $x \in (0, \pi/2)$**

1991 *Mathematics Subject Classification.* 41-XX,97-XX.

Key words and phrases. Inequality, trigonometry, polynomials, sin, cos, tan .

For $x \in (0, \pi/2)$ the inequality $x < \tan x$ yields $x \cos x < \sin x$, so for any $x \in (0, \pi/2)$ we have

$$x \cos x < \sin x < x \quad (5)$$

Using (5) we can find a third-degree polynomial that is a lower bound for $\sin x$. Indeed, since $\cos x = 1 - 2 \sin^2(x/2)$ and $\sin(x/2) < x/2$

we have $\cos x > 1 - 2 \left(\frac{x}{2}\right)^2 = 1 - \frac{x^2}{2}$ (which is the left side of (2)) and, therefore,

$$\sin x > x \cos x > x \left(1 - \frac{x^2}{2}\right) = x - \frac{x^3}{2}.$$

Suppose now that for some positive a the inequality $\sin x \geq x - ax^3$ holds for every $x \in (0, \pi/2)$. (We have just shown that this holds when $a = 1/2$).

Then, since $\cos x > 1 - x^2/2$, we obtain

$$\begin{aligned} \sin x &= 2 \sin \frac{x}{2} \cos \frac{x}{2} > 2 \left(\frac{x}{2} - a \left(\frac{x}{2}\right)^3\right) \left(1 - \frac{1}{2} \left(\frac{x}{2}\right)^2\right) = \\ &= x - \left(\frac{a}{4} + \frac{1}{8}\right) x^3 + \frac{1}{32} ax^5 > x - \left(\frac{a}{4} + \frac{1}{8}\right) x^3 \end{aligned}$$

whenever $x \in (0, \pi)$ and, therefore, for any $x \in (0, \pi/2)$ holds inequality

$$\sin x > x - \left(\frac{a}{4} + \frac{1}{8}\right) x^3 \iff x - \sin x < \left(\frac{a}{4} + \frac{1}{8}\right) x^3$$

So, we can see that if for some $a > 0$ inequality $x - \sin x < ax^3$ holds for any $x \in (0, \pi/2)$ then inequality $x - \sin x < \left(\frac{a}{4} + \frac{1}{8}\right) x^3$ holds for any $x \in (0, \pi/2)$ as well.

(Furthermore, even non-strict inequality $x - \sin x \leq ax^3$, $x \in (0, \pi/2)$ by procedure represented above generate strict inequality $x - \sin x < \left(\frac{a}{4} + \frac{1}{8}\right) x^3$ for $x \in (0, \pi)$).

Thus, for any term a_n of the sequence defined by $a_{n+1} = \frac{a_n}{4} + \frac{1}{8}$, $a_1 = 1/2$ inequality $r(x) < a_n x^3$ holds for any $x \in (0, \pi/2)$.

Since $\frac{1}{8} = c - \frac{c}{4} \iff c = \frac{1}{6}$ then for any $n \in \mathbb{N}$ we have

$$\begin{aligned} a_{n+1} &= \frac{a_n}{4} + \frac{1}{8} \iff a_{n+1} - \frac{1}{6} = \frac{1}{4} \left(a_n - \frac{1}{6}\right) \iff 4^{n+1} \left(a_{n+1} - \frac{1}{6}\right) = \\ &= 4^n \left(a_n - \frac{1}{6}\right). \end{aligned}$$

That is $4^n \left(a_n - \frac{1}{6}\right) = 4^1 \left(a_1 - \frac{1}{6}\right) = 4 \left(\frac{1}{2} - \frac{1}{6}\right) = \frac{4}{3}$ and, therefore, $a_n = \frac{1}{6} + \frac{1}{3 \cdot 4^{n-1}}$, $n \in \mathbb{N}$.

Thus, $x - \sin x - \frac{x^3}{6} < \frac{x^3}{3 \cdot 4^{n-1}} < \frac{\pi^3}{6 \cdot 4^n}$, $n \in \mathbb{N}$. Since $\frac{1}{4^n} < \frac{1}{3n}$ for any $n \in \mathbb{N}$ (can be proved by Math Induction) then $x - \sin x - \frac{x^3}{6} < \frac{\pi^3}{18n}$, $n \in \mathbb{N}$ and applying

Proposition to $f(x) = x - \sin x - \frac{x^3}{6}$ and $P = \left\{ \frac{\pi^3}{18n} \mid n \in \mathbb{N} \right\}$ we obtain inequality

$$x - \sin x - \frac{x^3}{6} \leq 0 \iff x - \sin x \leq \frac{x^3}{6} \text{ for any } x \in (0, \pi/2).$$

Since $x - \sin x \leq \frac{x^3}{6}$ yields $x - \sin x < \left(\frac{1}{6} \cdot \frac{1}{4} + \frac{1}{8} \right) x^3 = x - \frac{x^3}{6}$ we finally get strict inequality $x - \sin x < \frac{x^3}{6} \iff x - \frac{x^3}{6} < \sin x$ for any $x \in (0, \pi/2)$.

• **Proof of** $\cos x < 1 - \frac{x^2}{2} + \frac{x^4}{24}, x \in (0, \pi/2)$.

Since $\sin x > x - \frac{x^3}{6} > 0$ ($x - \frac{x^3}{6} = \frac{x}{24}(24 - 4x^2) > \frac{x}{24}(24 - \pi^2) > 0$), we have

$$\cos x = 1 - 2 \sin^2 \left(\frac{x}{2} \right) \leq 1 - 2 \left(\frac{x}{2} - \frac{1}{6} \left(\frac{x}{2} \right)^3 \right)^2 = 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{1152}x^6 < 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4.$$

• **Proof of** $\sin x < x - \frac{x^3}{3!} + \frac{x^5}{5!}, x \in (0, \pi/2)$

Let $x \in (0, \pi/2)$. Since $\sin x < x$ and $\cos x < 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4$ then

$$\sin x = 2 \sin \frac{x}{2} \cos \frac{x}{2} < 2 \cdot \frac{x}{2} \left(1 - \frac{(x/2)^2}{2} + \frac{(x/2)^4}{24} \right) = x - \frac{x^3}{8} + \frac{x^5}{384}.$$

So, we have

$$\sin x < x - \frac{x^3}{8} + \frac{x^5}{384}, \quad x \in (0, \pi/2). \quad (6)$$

Suppose now that for some positive $a < \frac{1}{3!} = \frac{1}{6}$ and positive

$b < \frac{1}{5!} = \frac{1}{120}$ inequality $\sin x \leq x - ax^3 + bx^5$ holds for every $x \in (0, \pi/2)$.

Then

$$\begin{aligned} \sin x &= 2 \sin \frac{x}{2} \cos \frac{x}{2} < 2 \left(\frac{x}{2} - a \left(\frac{x}{2} \right)^3 + b \left(\frac{x}{2} \right)^5 \right) \left(1 - \frac{1}{2} \left(\frac{x}{2} \right)^2 + \frac{1}{24} \left(\frac{x}{2} \right)^4 \right) = \\ &x - \left(\frac{a}{4} + \frac{1}{8} \right) x^3 + \left(\frac{a}{32} + \frac{b}{16} + \frac{1}{384} \right) x^5 - \left(\frac{x}{2} \right)^7 \left(\frac{1}{12} a + b \left(1 - \frac{x^2}{48} \right) \right) < \\ &x - \left(\frac{a}{4} + \frac{1}{8} \right) x^3 + \left(\frac{a}{32} + \frac{b}{16} + \frac{1}{384} \right) x^5 \end{aligned}$$

because for any $x \in (0, \pi/2)$ holds inequality $\frac{1}{12}a + b \left(1 - \frac{x^2}{48} \right) > 0$.

Thus, the non-strict inequality $\sin x \leq x - ax^3 + bx^5$ (which we know to be true when $a = 1/8$ and $b = 1/384$) yields the strict inequality

$$\sin x < x - \left(\frac{a}{4} + \frac{1}{8} \right) x^3 + \left(\frac{a}{32} + \frac{b}{16} + \frac{1}{384} \right) x^5 \quad (7)$$

If we write $a = \frac{1}{6} - p, b = \frac{1}{120} - q$ and denote $\sin x - x + \frac{x^3}{6} - \frac{x^5}{120}$

via $r(x)$ then

$$\sin x \leq x - ax^3 + bx^5 \iff \sin x \leq x - \left(\frac{1}{6} - p \right) x^3 + \left(\frac{1}{120} - q \right) x^5 \iff$$

$$\sin x - x + \frac{x^3}{6} - \frac{x^5}{120} \leq px^3 - qx^5 \iff r(x) \leq px^3 - qx^5, \text{ where } p, q > 0$$

and since

$$\begin{aligned} \frac{a}{4} + \frac{1}{8} &= \frac{1}{6} - \frac{p}{4}, \quad \frac{a}{32} + \frac{b}{16} + \frac{1}{384} = \\ \frac{1}{32} \left(\frac{1}{6} - p \right) + \frac{1}{16} \left(\frac{1}{120} - q \right) + \frac{1}{384} &= \frac{1}{120} - \frac{p}{32} - \frac{q}{16} \end{aligned}$$

$$\text{we obtain (7)} \iff \sin x < x - \left(\frac{1}{6} - \frac{p}{4} \right) x^3 + \left(\frac{1}{120} - \frac{p}{32} - \frac{q}{16} \right) x^5 \iff$$

$$r(x) < \frac{p}{4} x^3 - \left(\frac{p}{32} + \frac{q}{16} \right) x^5.$$

So, we have shown that the inequality $r(x) < px^3 - qx^5$ and even $r(x) \leq px^3 - qx^5$ implies the inequality

$$r(x) < \frac{p}{4} x^3 - \left(\frac{p}{32} + \frac{q}{16} \right) x^5.$$

Due to inequality (6) with $a = \frac{1}{8}$ and $b = \frac{1}{384}$ the initial value

of p is $\frac{1}{6} - \frac{1}{8} = \frac{1}{24}$ and the initial value of q is $\frac{1}{120} - \frac{1}{384} = \frac{11}{1920}$.

Let sequences $(p_n)_{n \geq 1}$ and $(q_n)_{n \geq 1}$ be as follows

$$p_{n+1} = \frac{p_n}{4}, \quad q_{n+1} = \frac{p_n}{32} + \frac{q_n}{16}, \quad n \in \mathbb{N}, \quad p_1 = \frac{1}{24}, \quad q_1 = \frac{11}{1920}.$$

Thus, we can see that for any $x \in (0, \pi/2)$ holds inequality

$$r(x) < p_n x^3 - q_n x^5 < p_n x^3 \quad (\text{because } q_n x^5 > 0 \text{ for any } n \in \mathbb{N})$$

Noting that $p_n = \frac{1}{24} \cdot \frac{1}{4^{n-1}} = \frac{1}{3 \cdot 2^{2n+1}}$, $n \in \mathbb{N}$ we obtain inequality

$$r(x) < p_n x^3 = \frac{x^3}{3 \cdot 2^{2n+1}}, \quad n \in \mathbb{N}.$$

Since $\frac{1}{3 \cdot 2^{2n+1}}$ can be arbitrary small with increasing n

then $\frac{x^3}{3 \cdot 2^{2n+1}} < \frac{(\pi/2)^3}{6 \cdot 4^n}$ can be arbitrary small with increasing n

as well then, applying **Proposition** to $f(x) = r(x)$ we obtain inequality $r(x) \leq 0$, $x \in (0, \pi/2)$ which equivalent to inequality

$$\sin x \leq x - \frac{1}{6} x^3 + \frac{1}{120} x^5, \quad x \in (0, \pi/2) \text{ and, since}$$

$$\sin x \leq x - \frac{1}{6} x^3 + \frac{1}{120} x^5 \text{ yields}$$

$$\sin x < x - \left(\frac{1}{6} \cdot \frac{1}{4} + \frac{1}{8} \right) x^3 + \left(\frac{1}{6} \cdot \frac{1}{32} + \frac{1}{120} \cdot \frac{1}{16} + \frac{1}{384} \right) x^5 =$$

$$x - \frac{1}{6} x^3 + \frac{1}{120} x^5$$

we finally get strict inequality $\sin x < x - \frac{1}{6} x^3 + \frac{1}{120} x^5$.

$$\bullet \text{ Proof of } 1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} < \cos x, \quad x \in (0, \pi/2).$$

As a consequence of the above inequalities for $\sin x$ we have

$$\cos x = 1 - 2 \sin^2 \left(\frac{x}{2} \right) \geq 1 - 2 \left(\frac{x}{2} - \frac{1}{3!} \left(\frac{x}{2} \right)^3 + \frac{1}{5!} \left(\frac{x}{2} \right)^5 \right)^2 =$$

$$1 - 2 \left(\frac{x^2}{4} - \frac{x^4}{3!2^3} + \frac{1}{45} \cdot \frac{x^6}{2^5} - \frac{1}{3!5!} \left(\frac{x}{2} \right)^8 \left(2 - \frac{1}{4 \cdot 5} \left(\frac{x}{2} \right)^2 \right) \right) =$$

$$1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + \frac{2}{3!5!} \left(\frac{x}{2} \right)^8 \left(2 - \frac{1}{4 \cdot 5} \left(\frac{x}{2} \right)^2 \right) > 1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720}$$

because $2\frac{1}{4 \cdot 5} \left(\frac{x}{2}\right)^2 < 2$ for $x \in (0, \pi/2)$.

• **Proof of** $x + x^3/3 < \tan x, x \in (0, \pi/2)$

Let $x \in (0, \pi/2)$. Since $\tan x > x$ we obtain

$$\tan x = \frac{2 \tan\left(\frac{x}{2}\right)}{1 - \tan^2\left(\frac{x}{2}\right)} > \frac{2 \cdot \frac{x}{2}}{1 - \left(\frac{x}{2}\right)^2} > x \left(1 + \frac{x^2}{4}\right) = x + \frac{x^3}{4}.$$

As above, suppose that for some $a > 0$ inequality $\tan x \geq x + ax^3$ holds for any $x \in (0, \pi/2)$. Then

$$\begin{aligned} \tan x &= \frac{2 \tan\left(\frac{x}{2}\right)}{1 - \tan^2\left(\frac{x}{2}\right)} > \frac{2 \left(\frac{x}{2} + \frac{ax^3}{8}\right)}{1 - \left(\frac{x}{2} + \frac{ax^3}{8}\right)^2} > \left(x + \frac{ax^3}{4}\right) \left(1 + \left(\frac{x}{2} + \frac{ax^3}{8}\right)^2\right) > \\ &\quad \left(x + \frac{ax^3}{4}\right) \left(1 + \frac{x^2}{4}\right) > x + \left(\frac{a}{4} + \frac{1}{4}\right)x^3. \end{aligned}$$

Thus, the non strict inequality $\tan x \geq x + ax^3$ (which we know to be true when $a = 1/4$) yields the strict inequality $\tan x > x + \left(\frac{a}{4} + \frac{1}{4}\right)x^3$.

If we write $a = \frac{1}{3} - p$, then $\frac{a}{4} + \frac{1}{4} = \frac{1}{3} - \frac{p}{4}$. So we have shown that the inequality $r(x) < px^3$ implies the inequality $r(x) < \frac{px^3}{4}$, where $r(x) = x + \frac{x^3}{3} - \tan x$.

We know that the inequality holds for $p = 1/12$ because $\tan x > x + \frac{x^3}{4} \iff r(x) < \frac{x^3}{12}$.

So, inequality $r(x) < px^3$ holds for $p = \frac{1}{48}, \frac{1}{192}, \dots, \frac{1}{12 \cdot 4^n}, \dots$ in fact for arbitrary small positive p . Therefore, applying **Proposition** to the function $f(x) = r(x)$ we obtain

$r(x) \leq 0 \iff x + \frac{x^3}{3} \leq \tan x$ for $x \in (0, \pi/2)$ and since $\tan x \geq x + \frac{1}{3}x^3$ yields $\tan x > x + \left(\frac{1}{3} \cdot \frac{1}{4} + \frac{1}{4}\right)x^3 = x + \frac{1}{3}x^3$ we finally get strict inequality $\tan x > x + \frac{1}{3}x^3$.