



## József Wildt International Mathematical Competition

The Edition XXVI<sup>th</sup>, 2016<sup>22</sup>

The solution of the problems W.1 - W.50 must be mailed before 10. September 2016, to Mihály Bencze, str. Hărmanului 6, 505600 Săcele - Négyfalu, Jud. Braşov, Romania, E-mail: [benczemihaly@yahoo.com](mailto:benczemihaly@yahoo.com); [benczemihaly@gmail.com](mailto:benczemihaly@gmail.com)

**W1.** Show that  $H = \sum_{\substack{q \leq Q \\ q \text{ odd, squarefree}}} \prod_{p|q} \frac{2}{p-2} \gg (\log Q)^2$ .

Michael Rassias

**W2.** Let  $D$  be a fixed squarefree integer with  $D \geq 1$ . Let also

$$\pi_D(x) = \sum_{\substack{p \leq x \\ \left(\frac{-D}{p}\right) = 1}} 1.$$

Prove that  $\pi_D(x) \sim \frac{1}{2}\pi(x)$ , when  $x \rightarrow +\infty$

Michael Rassias

**W3.** Let  $(F_k)_{k \geq 0}$ ,  $F_0 = 0$ ,  $F_1 = 1$ ,  $F_{k+2} = F_k + F_{k+1}$ ,  $(\forall) k \in N$  and  $A = (a_{ij})_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}}$ ,  $B = (b_{jk})_{\substack{1 \leq j \leq n \\ 1 \leq k \leq m}}$ ,  $C = (c_{rs})_{1 \leq r, s \leq m}$ ,  $a_{ij} = F_j$ ,  $b_{jk} = F_j$ ,  $c_{rs} = F_{m-r+1}^2$ ,  $(\forall) i, k = \overline{1, m}$ ,  $(\forall) \overline{1, n}$ . For  $p, q$  positive integers compute  $(AB)^p C^q$ .

D.M.Bătineţu-Giurgiu and Neculai Stanciu

**W4.** Let  $a, b \in (-\infty, +\infty)$ . Find  $\lim_{n \rightarrow \infty} \left( \frac{(n+3)^{n+2+a}}{(n+2)^{n+1+b}} - \frac{(n+2)^{n+1+a}}{(n+1)^{n+b}} \right)$ .

D.M.Bătineţu-Giurgiu and Neculai Stanciu

<sup>22</sup>Received 15.03.2016

2000 *Mathematics Subject Classification.* 11-06.

*Key words and phrases.* Contest.

**W5.** Let  $f : R_+^* \rightarrow R_+^*$  be a continue function such that  $\lim_{x \rightarrow \infty} \frac{f(x)}{x^m} = a \in R_+^*$ , where  $m \in [1, +\infty)$ . Calculate  $\lim_{n \rightarrow \infty} \left( \sqrt[n+1]{\prod_{k=1}^{n+1} f(k)} - \sqrt[n]{\prod_{k=1}^n f(k)} \right)$ .

D.M.Bătinețu-Giurgiu and Neculai Stanciu

**W6.** If  $a, b, c$  are the sides of a triangle, demonstrate the inequality  $\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} + \frac{r}{R} \leq 2$ .

Stănescu Florin

**W7.** If all triangle  $ABC$  holds

$$\sum \sin A - \prod \sin A \geq \sum \sin^3 A \geq \prod \sin A + 4 \prod \cos A (\sum \sin A).$$

Stănescu Florin

**W8.** Let  $f, g : [0, 1] \rightarrow (0, +\infty)$ ,  $f(0) = g(0) = 0$  two continuous functions such that  $f$  it's convex, and  $g$  concave. If  $h : [0, 1] \rightarrow R$  is an increasing function, show that  $\int_0^1 h(x) g(x) dx \int_0^1 f(x) dx \leq \int_0^1 g(x) dx \int_0^1 h(x) f(x) dx$ .

Stănescu Florin

**W9.** Let  $n$  be a positive integer. Prove that

$$\sum_{k=1}^n \frac{F_{2k}}{(F_{2k+1} - 1)^2} < 2,$$

where  $F_n$  is the  $n^{th}$  Fibonacci number. That is,  $F_0 = 0, F_1 = 1$  and  $F_{n+2} = F_{n+1} + F_n$  for  $n \geq 1$ .

José Luis Díaz-Barrero

**W10.** Let  $a, b$ , and  $c$  be positive real numbers. Prove that

$$\left( \frac{(6n+1)a-b}{n(b+c)} \right)^2 + \left( \frac{(6n+1)b-c}{n(c+a)} \right)^2 + \left( \frac{(6n+1)c-a}{n(a+b)} \right)^2 \geq 27$$

for any positive integer  $n \geq 1$ .

José Luis Díaz-Barrero

**W11.** Let  $x > y > z > t$  be four positive integers such that

$$(x^2 - y^2) + (xz - yt) - (z^2 - t^2) = 0.$$

Prove that  $xy + zt$  is a composite number.

José Luis Díaz-Barrero

**W12.** Let  $n \in \mathbb{N}$  and let  $O_n = 1 + \frac{1}{3} + \dots + \frac{1}{2n-1}$ . Calculate

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left(1 + \frac{2O_n}{n}\right)^n.$$

Ovidiu Furdui

**W13.** Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence of real numbers such that

$\lim_{n \rightarrow \infty} n(a_n - 1) = l \in (-\infty, \infty)$  and let  $p \geq 1$  be a natural number. Calculate

$$\lim_{n \rightarrow \infty} \prod_{k=1}^n \left(a_n + \frac{1}{\sqrt[k]{kn}}\right).$$

Ovidiu Furdui

**W14.** 1). Let  $f, g : [a, b] \rightarrow \mathbb{R}$  be two nonnegative continuous functions.

Assume that  $f$  attains its maximum at a unique point on  $[a, b]$  and  $g$  attains its maximum at the same point as  $f$  and possibly at other points. Prove that

$$\lim_{n \rightarrow \infty} \frac{\int_a^b f^{n+1}(x)g(x)dx}{\int_a^b f^n(x)dx} = \|f\|_\infty \|g\|_\infty$$

2). Does the result hold under no assumption on  $f$  and  $g$ ?

Ovidiu Furdui

**W15.** Let  $a, b, c$  be positive real numbers. Prove that

$$\sum_{\text{cyc}} \left( \frac{2a^2}{a+b} + \frac{a^3}{b^2+c^2} \right) \geq \frac{9}{2} \frac{a^2+b^2+c^2}{a+b+c}$$

Paolo Perfetti

**W16.** Prove that  $\sin t \left( \cos^3 t + \sqrt{\sin t} \right) \geq \left( \frac{2t}{\pi} \right)^{\cos^2 t}$  when  $0 \leq t \leq \frac{\pi}{2}$ .

Paolo Perfetti

**W17.** Let  $p$  a positive real number and let  $\{a_n\}_{n \geq 1}$  be a sequence defined by  $a_1 = 1, a_{n+1} = \frac{a_n}{1+a_n^p}$ . Find those real values  $q \neq 0$  such that the following series converges  $\sum_{n=1}^{\infty} \left| (pn)^{-\frac{1}{p}} - a_n \right|^q$ .

Paolo Perfetti

**W18.** In 3-dimensional Euclidean space, let  $S$  be a sphere with centre  $O$  and radius  $R$ . Three pairwise orthogonal rays originating in  $O$  intersect  $S$  at  $A, B, C$ . Let  $P$  be a point of  $S$  and let  $a, b, c$  denote the areas of triangles  $OPA, OPB, OPC$ , respectively. Prove  $2a(+b+c)(a^3+b^3+c^3) \geq R^4(ab+bc+ca)$ .

Michael Battaille

**W19.** Let  $a, b, c, x, y, z$  be real numbers such that  $a, b, c > 0$  and  $a+b+c = x+y+z = 1$ . Let  $u = xy - bz, v = az - cx, w = bx - ay$  and  $M = \begin{pmatrix} \frac{2y}{b+c} & z-c & b-y \\ c-z & \frac{2v}{c+a} & x-a \\ y-b & a-x & \frac{2w}{a+b} \end{pmatrix}$ . Prove that  $\det(M) = 0$  if and only if  $u+v+w = 0$ .

Michael Battaille

**W20.** Let  $\Gamma$  be a circle and let  $A \in \Gamma, B \in \Gamma, C \notin \Gamma$  be such that  $CA$  and  $CB$  intersect  $\Gamma$  again at diametrically opposite points. If  $l$  is the radial axis of  $\Gamma$  and the circle with centre  $A$ , radius  $AC$ , show that  $d(C, l) = \frac{CA \cdot CB}{AB}$ . ( $d(C, l)$  denotes the distance from  $C$  to  $l$ ).

Michael Battaille

**W21.** Let  $ABC$  be a triangle and we note  $AB = c, BC = a, CA = b$  and  $m_a, m_b, m_c$  the medians lengths corresponding to the vertexes  $A, B$  respective  $C$ . Prove the inequality  $\sum \frac{(b^2+c^2)^2}{m_a^3} \geq \frac{32\sqrt{3}}{9}(a+b+c)$ .

Ovidiu Pop

**W22.** Let  $ABCD$  be a convex quadrilateral,  $O \in [AC], OM \parallel BC, M \in AB, ON \parallel AB, N \in BC, OP \parallel AD, P \in CD$  and  $OQ \parallel CD, Q \in DA$ . Prove the inequalities  $\min \{Area[ABD], Area[BCD]\} \leq Area[MNPQ] \leq$

$$\leq \max \{ \text{Area}[ABD], \text{Area}[BCD] \}.$$

Ovidiu Pop

**W23.** Let  $n \in \mathbb{N}$ , for  $k$  integer,  $1 \leq k \leq n$ , euclidean division  $n$  by  $k$  gives  $n = qk + n_k$ , and denote  $p_n$  the probability that  $n_k \geq \frac{k}{2}$ . Calculate  $p_n$  and find  $\lim_{n \rightarrow \infty} p_n$ .

Moubinol Omarjee

**W24.** Let  $f \in C^3(\mathbb{R}^n, \mathbb{R})$  with  $f(0) = f'(0) = 0$ . Prove that there exist  $h \in C^3(\mathbb{R}^m, S_n(\mathbb{R}))$ , such that  $f(x) = x^t h(x) x$ , when  $S_n(\mathbb{R})$ , is the set of symmetric matrix, and  $x^t$  is the transpose of  $x$ .

Moubinol Omarjee

**W25.** Find the nature of the series  $\sum_{n \geq 1} \frac{e^{i \ln(p_n)}}{p_n}$  when  $(p_n)_{n \geq 1}$  is the prime number increasing order, and  $i$  imaginary complex number.

Moubinol Omarjee

**W26.** Let  $M$  be a point in the interior of triangle  $ABC$  and  $R_a, R_b, R_c$  the radii of circumcircle of  $MBC, MCA, MAB$ . Show that  $\frac{1}{R_a} + \frac{1}{R_b} + \frac{1}{R_c} \leq \frac{1}{MA} + \frac{1}{MB} + \frac{1}{MC}$

Nicuşor Minculete

**W27.** Let  $a_j > 0$ , ( $j = 1, 2, \dots, k$ ) such that  $\sum_{cyclic} \prod_{j=1}^{k-1} a_j = k$ , and  $n > 1$ .

Prove that  $\sum_{cyclic} \sqrt[n]{a_1 + \frac{1}{\prod_{j=1}^k a_j}} \geq k \sqrt[n]{2}$ .

Ángel Plaza

**W28.** Let  $(x_n)_{n \geq 0}$  be the sequence defined recurrently by

$x_{n+2} = x_{n+1} - \frac{1}{2}x_n$  with initial terms  $x_0 = 2$  and  $x_1 = 1$ . Find  $\sum_{n=1}^{\infty} \frac{x_n}{n+2}$ .

Ángel Plaza

**W29.** Let  $x, y, z$  be positive real numbers such that  $x + y + z = 1$ . Prove

$$\text{that } \sqrt{\frac{x^3 + 1}{x^2 + y + z}} + \sqrt{\frac{y^3 + 1}{y^2 + z + x}} + \sqrt{\frac{z^3 + 1}{z^2 + x + y}} \leq 3\sqrt{2}.$$

Ángel Plaza

**W30.** Let be  $x_0 = x_1 = 1$  and  $x_{n+2} = 5x_{n+1} - x_n - 1$  for all  $n \geq 0$ . Prove  $x_n x_m x_p x_{m+1} x_{m+2} x_{p+2} \geq \left( \sqrt[3]{x_{n+1}^2 x_{m+1}} + \sqrt[3]{x_{m+1}^2 x_{p+1}} + \sqrt[3]{x_{p+1}^2 x_{n+1}} \right)^3$  for all  $n, m, p \in N$ .

Mihály Bencze

**W31.** If  $a_k > 1$  ( $k = 1, 2, \dots, n$ ) then  $\sum (\log_{a_1} a_1 a_2)^\lambda \geq n2^\lambda$  for all  $\lambda \in (-\infty, 0] \cup [1, +\infty)$ .

Mihály Bencze

**W32.** If  $x, y, z > 0$  and  $x + y + z = 1$  then  $8(1-x)^x (1-y)^y (1-z)^z \leq 3(2-x-y)^{x+y} (2-y-z)^{y+z} (2-z-x)^{z+x}$ .

Mihály Bencze

**W33.** 1). Let  $f : R \rightarrow R$  be a bijective and continuous function such that  $f(a) = \lambda a$  and  $f(b) = \lambda b$  when  $a, b \in R$  and  $\lambda \in R^*$ . Prove that exist  $x_0 \in (a, b)$  such that  $f(x_0) + \lambda f^{-1}(\lambda x_0) = 2\lambda x_0$ .

2). Let  $f : R \rightarrow R$  be a bijective and continuous function such that  $f(a) = \lambda b$  and  $f(b) = \lambda a$  when  $a, b \in R$  and  $\lambda \in R^*$ . Prove that exist  $x_0 \in (a, b)$  such that  $f\left(\frac{f(x_0)}{\lambda}\right) = \lambda x_0$ .

Mihály Bencze

**W34.** Prove  $\sum_{k=1}^n \left( \frac{2(k+1)(k+2)^2}{((k+2)!)^3} \right)^{\frac{1}{k}} \geq \frac{n(n+5)}{3(n+2)(n+3)}$ .

Mihály Bencze

**W35.** Prove that exist infinitely many  $n \in N$  for which  $n!$  is divisible by  $n^5 + n - 1$ .

Mihály Bencze

**W36.** Let  $\Delta(x, y, z) = 2(xy + yz + zx) - (x^2 + y^2 + z^2)$  and let  $a, b, c$  be sidelengths of a triangle with area  $F$ . Prove that  $\Delta(a^3, b^3, c^3) \leq \frac{64F^3}{\sqrt{3}}$ .

Arkady Alt

**W37.** Let  $E$  be an inner Product Space with dot product  $\cdot$  and  $F$  be proper nonzero subspace. Let  $P : E \rightarrow E$  be orthogonal projection of  $E$  on  $F$ .

a). Prove that for any  $x, y \in E$ , holds inequality

$$|x \cdot y - xP(y) - yP(x)| \leq \|x\| \cdot \|y\|$$

b). Determine all cases when equality occurs

Arkady Alt

**W38.** Prove that  $0 < \left(\frac{4^x + 2^x + 1}{x}\right)^x - 2^x < 1$  for all  $x \in \left(0, \frac{1}{2e}\right]$ .

Ionel Tudor

**W39.** Let  $n \geq 2$  be a natural number and  $a_i > 0, i = \overline{1, n}$ . If  $S = \sum_{i=1}^n a_i$  and  $x_i = S - a_i$ , then the following inequality holds:

$$\frac{\prod_{i=1}^n \sqrt{a_i}}{\sqrt[n-1]{\prod_{1 \leq i < j \leq n} (a_i + a_j)}} \leq \frac{\prod_{i=1}^n \sqrt{x_i}}{\sqrt[n-1]{\prod_{1 \leq i < j \leq n} (x_i + x_j)}}.$$

Ovidiu Bagdasar

**W40.** Prove that if  $x_i > 0, i = \overline{1, n}$ , then the next inequality holds:

$$\sum_{i=1}^n \frac{S_{\alpha+\beta} - x_i^{\alpha+\beta}}{S_{\alpha} - x_i^{\alpha}} \leq n \cdot \frac{S_{\alpha+\beta}}{S_{\alpha}}, \quad (33)$$

provided that  $\alpha\beta \geq 0$  and  $S_p = \sum_{i=1}^n x_i^p$ , for any real number  $p$ .

Ovidiu Bagdasar

**W41.** Let  $n \geq 2$  a natural number and the numbers  $a_i > 1, i = \overline{1, n}$ . Prove that

$$\sum_{i=1}^n \frac{\log_{a_i} a_{i+1}^{n-1}}{S - a_i} \geq \frac{n^2}{\sum_{i=1}^n a_i}.$$

Ovidiu Bagdasar

**W42.** Let  $ABC$  be an acute triangle. The angle bisectors from  $A, B, C$  meet the opposite sides in  $A_1, B_1, C_1$ , respectively. Let  $R$  and  $r$  be the circumradius and the inradius of the triangle  $ABC$ , respectively. Let  $R_A, R_B$ , and  $R_C$  the circumradii of the triangles  $AC_1B_1, BA_1C_1$ , and  $CB_1A_1$ , respectively. Prove that

$$R_A + R_B + R_C \geq R + r.$$

Pál Péter Dályay

**W43.** Let  $f$  be a continuous real function defined on the set of the nonnegative real numbers for which the following integrals are convergent:  $S = \int_0^\infty f^2(x)dx, T = \int_0^\infty x f^2(x)dx, U = \int_0^\infty x^2 f^2(x)dx$ . Prove that

$$\left( \int_0^\infty |f(x)|dx \right)^2 \leq 2 \left( \sqrt{SU} + T \right).$$

Pál Péter Dályay

**W44.** If  $\zeta$  is the Riemann zeta-function, and  $s$  is a real number greater than  $3/2$ , then:

$$\begin{aligned} \zeta^2(s) &\leq \pi\sqrt{2} \left( \sum_{i,j=1}^{\infty} \frac{1}{i^{s-1}j^{s-1}(i+j)} \sum_{k,l=1}^{\infty} \frac{1}{k^{s-1}l^{s-1}(k+l)^3} \right)^{1/2} \leq \\ &\leq \frac{\pi}{2\sqrt{2}} \zeta(s-1/2)\zeta(s+1/2). \end{aligned}$$

Pál Péter Dályay

**W45.** Let  $x \in \text{Poisson}(2)$  be a random variable

i). Find the set  $M$  of all the values  $n \in \mathbb{N}^*$  so that

$$P\left(\left\{ \omega / |x(\omega) - 2| \geq \frac{2}{n} \right\}\right) \leq \frac{128}{n^2}$$

ii). From the set  $M$  we extract 2 numbers one, after the other. Find the probability that the second extracted number could be the greatest, in the following two situation: with return in the set, and without return in  $M$ .

Laurențiu Modan



- W46.** i). For any  $n > 3$  natural numbers, prove that  $2n! + (2n)! > 3 \cdot 2^{n+2}$   
 ii). Study the convergence of the series  $\sum_{n>3} \frac{1}{2n!+(2n)!}$

Laurențiu Modan

**W47.** Let  $a, b, c, d \in (0, 1)$  and  $f : (0, 1) \rightarrow R$  a convex and decreasing function. Prove that  $\sum f(1 - a^3) \geq \sum f(1 - a^2b)$ .

Marius Drăgan

**W48.** Let  $I$  be an interval and  $f : I \rightarrow R$  a convex function and

$x_1, x_2, \dots, x_n \in I$ . Prove that  $\sum_{k=1}^n f(x_k) - nf\left(\frac{1}{n} \sum_{k=1}^n x_k\right) \geq$   
 $\max_{1 \leq i < \dots < i_k \leq n} \left( f(x_{i_1}) + \dots + f(x_{i_k}) - kf\left(\frac{x_{i_1} + \dots + x_{i_k}}{k}\right) \right)$

Marius Drăgan and Mihály Bencze

**W49.** Find all the functions  $f : R \rightarrow [-1, 1]$  such that  $f(x+y)(1 - f(x)f(y)) = f(x) + f(y)$  for each  $x, y \in R$ .

Marius Drăgan and Sorin Rădulescu

**W50.** Find all the function  $f : R \rightarrow R$  which are continuous in a real point  $x_0$  such that  $f(x+y) = f(x)\sqrt{1+f^2(y)} + f(y)\sqrt{1+f^2(x)}$  for each  $x, y \in R$ .

Sorin Rădulescu